

**CORRIGENDUM TO “ L^p -GENERIC COCYCLES HAVE
ONE-POINT LYAPUNOV SPECTRUM”**

ALEXANDER ARBIETO AND JAIRO BOCHI

Lemma 4 in [1] is not true, unless $k = 1$ (this case was used in page 78, line 9). For arbitrary k , the correct statement is:

Lemma 4’. *Given $k \in \{1, \dots, d\}$ and $P > 0$, there is $C = C(P, k, d) > 0$ such that for every $A, B \in \text{GL}(d, \mathbb{R})$ with $\|A\|, \|B\| \leq P$, we have*

$$\log^+ \|\wedge^k B\| \leq \log^+ \|\wedge^k A\| + C\|B - A\|.$$

Proof. Let $L = \wedge^k B - \wedge^k A$. Let $\{e_i; 1 \leq i \leq d\}$ be an orthonormal basis of \mathbb{R}^d . Then $\{e_{i_1} \wedge \dots \wedge e_{i_k}; 1 \leq i_1 < \dots < i_k \leq d\}$ is an orthonormal basis of $\wedge^k \mathbb{R}^d$. We have

$$\begin{aligned} L(e_{i_1} \wedge \dots \wedge e_{i_k}) &= (B - A)e_{i_1} \wedge Be_{i_2} \wedge \dots \wedge Be_{i_k} + \\ &\quad Ae_{i_1} \wedge (B - A)e_{i_2} \wedge Be_{i_3} \wedge \dots \wedge Be_{i_k} + \dots + \\ &\quad Ae_{i_1} \wedge \dots \wedge Ae_{i_{k-1}} \wedge (B - A)e_{i_k}. \end{aligned}$$

It follows that $\|L(e_{i_1} \wedge \dots \wedge e_{i_k})\| \leq kP^{k-1}\|B - A\|$. Therefore $\|L\| \leq C\|B - A\|$ for some constant C . Then $\|\wedge^k B\| \leq \|\wedge^k A\| + C\|B - A\|$ and using that $\log^+(x + y) \leq \log^+ x + y$ for all $x, y \geq 0$, the lemma follows. \square

Now we indicate the necessary modifications in the proof of part (a) of Theorem 2. In the case condition (3) is satisfied, we let N and K be as before. Take $P = e^{KN}$ and let C be given by lemma 4’. We modify the definition of δ' in (5) to $\delta' = \min\{\eta, \varepsilon C^{-1}e^{-K(N-1)}\}$. If $x \in G$ then $\|A^N(x)\|, \|B^N(x)\| \leq P$. So we can apply lemma 4’ and the estimative (8) becomes

$$\frac{1}{N} \int_G \log^+ \|\wedge^k B^N\| \leq \Lambda_k(A) + 2\varepsilon + \frac{C}{N} \int_G \|B^N - A^N\|.$$

Then the upper bound in (9) becomes $\Lambda_k(A) + 3\varepsilon$, and so the first case of part (a) follows.

For the general case, we replace the first part in (10) with $k \int_{L_a} \log^+ \|A\| < \varepsilon$. Then in page 80, line 2, we can further estimate

$$\dots \leq \int_{L_a} \log^+ \|\wedge^k B\| \leq k \int_{L_a} \log^+ \|B\|.$$

Using lemma 4 with $k = 1$, we can change line 4 to

$$\int_{L_a} \hat{\lambda}_k(B, x) d\mu(x) \leq k \int_{L_a} \log^+ \|A\| + k \int_{L_a} \|B - A\|.$$

Then we finish the proof in the same way as before.

Date: July 15, 2003.

REFERENCES

- [1] A. ARBIETO AND J. BOCHI. L^p -generic cocycles have one-point Lyapunov spectrum. *Stochastics and Dynamics*, 3 (2003) no. 1, 73–81.

Alexander Arbieto (alexande@impa.br)

Jairo Bochi (bochi@impa.br)

IMPA, Estrada D. Castorina 110, Jardim Botânico, 22460-320 Rio de Janeiro, Brazil