

Generic Symplectic Diffeomorphisms and Partial Hyperbolicity

Workshop on Symplectic Dynamics
Institute for Advanced Study (Princeton)

Jairo Bochi

Pontifícia Universidade Católica do Rio de Janeiro

Plan of the talk

- Recall basic definitions.
- State a theorem on partial hyperbolicity of generic symplectic diffeomorphisms.
- Discuss consequences and further developments of that theorem (including the ergodicity result).
- Compare with a cousin theorem for volume-preserving diffeomorphisms.
- Sketch the main ideas of the proof, and so explain:
 - ▶ why symplectic is more difficult than volume-preserving;
 - ▶ the probabilistic method for constructing the perturbations.

Lyapunov exponents, Oseledets splitting

$f : M \rightarrow M$ diffeomorphism of a compact manifold of dimension d .
By the **Oseledets theorem**, there exists a full probability set $R \subset M$ such that for every (*regular point*) $x \in R$ there is a (*Oseledets*) splitting

$$T_x M = E^1(x) \oplus \cdots \oplus E^{k(x)}(x), \quad (\text{each } \neq \{0\})$$

and numbers (*Lyapunov exponents*) $\lambda_1(x) > \cdots > \lambda_{k(x)}(x)$ such that

$$\frac{1}{n} \log \|Df^n(x) \cdot v\| \xrightarrow{n \rightarrow \pm\infty} \lambda_i(x) \quad \forall v \in E^i(x) \setminus \{0\}.$$

Lyapunov exponents, Oseledets splitting

$f : M \rightarrow M$ diffeomorphism of a compact manifold of dimension d .
By the **Oseledets theorem**, there exists a full probability set $R \subset M$ such that for every (*regular point*) $x \in R$ there is a (*Oseledets*) splitting

$$T_x M = E^1(x) \oplus \cdots \oplus E^{k(x)}(x), \quad (\text{each } \neq \{0\})$$

and numbers (*Lyapunov exponents*) $\lambda_1(x) > \cdots > \lambda_{k(x)}(x)$ such that

$$\frac{1}{n} \log \|Df^n(x) \cdot v\| \xrightarrow{n \rightarrow \pm\infty} \lambda_i(x) \quad \forall v \in E^i(x) \setminus \{0\}.$$

The *zipped Oseledets splitting* is obtained by summing together all spaces with exponents of the same sign:

$$T_x M = E^+(x) \oplus E^0(x) \oplus E^-(x).$$

($E^*(x) = \{0\}$ now allowed, of course.)

Lyapunov exponents, Oseledets splitting (continued)

The multiplicity of each Lyapunov exponent $\lambda_j(x)$ is $\dim E^j(x)$ (by definition).

Indicate the Lyapunov exponents repeated according to multiplicity by:

$$\lambda_1(x) \geq \lambda_2(x) \geq \dots \geq \lambda_d(x), \quad (d = \dim M).$$

If f preserves a **symplectic** form ω on M then (d is even and) the exponents are symmetric:

$$\lambda_1 = -\lambda_d, \quad \lambda_2 = -\lambda_{d-1}, \quad \dots, \quad \lambda_{\frac{d}{2}} = -\lambda_{\frac{d}{2}+1}.$$

In particular, if $T_x M = E^+(x) \oplus E^0(x) \oplus E^-(x)$ is the zipped Oseledets splitting then

$$\dim E^+(x) = \dim E^-(x), \quad \dim E^0(x) = \text{even}.$$

A quote

Mañé ICM lecture (1983):

*Oseledets' theorem is essentially a measure theoretical result and therefore the information it provides holds only in that category. For instance, the **Lyapunov splitting is just a measurable function of the point***

A quote

Mañé ICM lecture (1983):

Oseledets' theorem is essentially a measure theoretical result and therefore the information it provides holds only in that category. For instance, the Lyapunov splitting is just a measurable function of the point and the limits defining the Lyapunov exponents are not uniform. It is clear that this is not a deficiency of the theorem but the natural counterweight to its remarkable generality.

A quote

Mañé ICM lecture (1983):

Oseledets' theorem is essentially a measure theoretical result and therefore the information it provides holds only in that category. For instance, the Lyapunov splitting is just a measurable function of the point and the limits defining the Lyapunov exponents are not uniform. It is clear that this is not a deficiency of the theorem but the natural counterweight to its remarkable generality. However, one can pose the problem (...) of whether these aspects can be substantially improved by working under generic conditions.

Invariant splittings with uniform properties

We'll consider Df -invariant splittings of $T_\Lambda M$, where Λ is a f -invariant set.

Invariant splittings with uniform properties

We'll consider Df -invariant splittings of $T_\Lambda M$, where Λ is a f -invariant set.

- 1 A splitting $T_\Lambda M = E^u \oplus E^s$ is called *uniformly hyperbolic* if E^s is uniformly contracted and E^u is uniformly expanded (i.e. contracted in the past): there are constants $C > 0$, $\tau > 1$ such that

$$\forall n \geq 0, \begin{cases} \|Df^n(x) \cdot v^s\| \leq C\tau^{-n}\|v^s\| & \forall v^s \in E^s(x) \setminus \{0\} \\ \|Df^{-n}(x) \cdot v^u\| \leq C\tau^{-n}\|v^u\| & \forall v^u \in E^u(x) \setminus \{0\} \end{cases}$$

Invariant splittings with uniform properties

We'll consider Df -invariant splittings of $T_\Lambda M$, where Λ is a f -invariant set.

- 1 A splitting $T_\Lambda M = E^u \oplus E^s$ is called *uniformly hyperbolic* if E^s is uniformly contracted and E^u is uniformly expanded (i.e. contracted in the past): there are constants $C > 0$, $\tau > 1$ such that

$$\forall n \geq 0, \begin{cases} \|Df^n(x) \cdot v^s\| \leq C\tau^{-n}\|v^s\| & \forall v^s \in E^s(x) \setminus \{0\} \\ \|Df^{-n}(x) \cdot v^u\| \leq C\tau^{-n}\|v^u\| & \forall v^u \in E^u(x) \setminus \{0\} \end{cases}$$

- 2 A splitting $T_\Lambda M = E^u \oplus E^c \oplus E^s$ is called *partially hyperbolic* if:
 - ▶ E^u is uniformly expanding (i.e. contracted in the past) and E^s is uniformly contracting;

Invariant splittings with uniform properties

We'll consider Df -invariant splittings of $T_\Lambda M$, where Λ is a f -invariant set.

- ① A splitting $T_\Lambda M = E^u \oplus E^s$ is called *uniformly hyperbolic* if E^s is uniformly contracted and E^u is uniformly expanded (i.e. contracted in the past): there are constants $C > 0$, $\tau > 1$ such that

$$\forall n \geq 0, \begin{cases} \|Df^n(x) \cdot v^s\| \leq C\tau^{-n}\|v^s\| & \forall v^s \in E^s(x) \setminus \{0\} \\ \|Df^{-n}(x) \cdot v^u\| \leq C\tau^{-n}\|v^u\| & \forall v^u \in E^u(x) \setminus \{0\} \end{cases}$$

- ② A splitting $T_\Lambda M = E^u \oplus E^c \oplus E^s$ is called *partially hyperbolic* if:
- ▶ E^u is uniformly expanding (i.e. contracted in the past) and E^s is uniformly contracting;
 - ▶ E^u **dominates** E^c , and E^c **dominates** E^s .

Invariant splittings with uniform properties

We'll consider Df -invariant splittings of $T_\Lambda M$, where Λ is a f -invariant set.

- 1 A splitting $T_\Lambda M = E^u \oplus E^s$ is called *uniformly hyperbolic* if E^s is uniformly contracted and E^u is uniformly expanded (i.e. contracted in the past): there are constants $C > 0$, $\tau > 1$ such that

$$\forall n \geq 0, \begin{cases} \|Df^n(x) \cdot v^s\| \leq C\tau^{-n}\|v^s\| & \forall v^s \in E^s(x) \setminus \{0\} \\ \|Df^{-n}(x) \cdot v^u\| \leq C\tau^{-n}\|v^u\| & \forall v^u \in E^u(x) \setminus \{0\} \end{cases}$$

- 2 A splitting $T_\Lambda M = E^u \oplus E^c \oplus E^s$ is called *partially hyperbolic* if:
 - ▶ E^u is uniformly expanding (i.e. contracted in the past) and E^s is uniformly contracting;
 - ▶ E^u **dominates** E^c , and E^c **dominates** E^s .

Remark: These bundles are automatically uniformly continuous, and thus extend (with the same hyperbolicity properties) to the closure $\overline{\Lambda}$.

Domination: the weakest uniform form of hyperbolicity

If E, F are Df -invariant subbundles of $T_\Lambda M$ then we say that E dominates F (in symbols, $E \succ F$), if there are constants $c > 0, \tau > 1$ such that for all unit vectors $\vec{e} \in E(x), \vec{f} \in F(x)$ and all $n \geq 0$

$$\frac{\|Df^n(x) \cdot \vec{e}\|}{\|Df^n(x) \cdot \vec{f}\|} > c\tau^n.$$


Domination: the weakest uniform form of hyperbolicity

If E, F are Df -invariant subbundles of $T_\Lambda M$ then we say that E *dominates* F (in symbols, $E \succ F$), if there are constants $c > 0, \tau > 1$ such that for all unit vectors $\vec{e} \in E(x), \vec{f} \in F(x)$ and all $n \geq 0$

$$\frac{\|Df^n(x) \cdot \vec{e}\|}{\|Df^n(x) \cdot \vec{f}\|} > c\tau^n.$$

Dominated splitting: $T_\Lambda M = E^1 \oplus \dots \oplus E^k$ with $E^1 \succ E^2 \succ \dots \succ E^k$.

“Morse–Smale-like” dynamics on projective space:



Domination is a.k.a. (in ODE theory) as *exponential separation*. It dates back to Perron (rediscovered by Mañé.)

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Theorem (B. '10)

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold:

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Theorem (B. '10)

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledets splitting $T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Theorem (B. '10)

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledets splitting $T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

- 1 either *trivial* $T_\Lambda M = E^0$, i.e., all Lyapunov exponents at x are zero;

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Theorem (B. '10)

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledec's splitting $T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

- 1 either *trivial* $T_\Lambda M = E^0$, i.e., all Lyapunov exponents at x are zero;
- 2 or *uniformly hyperbolic* with $E^u = E^+$, $E^s = E^-$;

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Theorem (B. '10)

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledec's splitting

$T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

- 1 either *trivial* $T_\Lambda M = E^0$, i.e., all Lyapunov exponents at x are zero;
- 2 or *uniformly hyperbolic* with $E^u = E^+$, $E^s = E^-$;
- 3 or *partially hyperbolic* with $E^u = E^+$, $E^c = E^0$, $E^s = E^-$ (all $\neq \{0\}$).

Mañé's statement

$\text{Diff}_\omega^1(M)$ is the set of symplectic C^1 -diffeomorphisms of (M, ω) , endowed with the C^1 topology. We consider on M the volume measure μ induced by ω .

Theorem (B. '10)

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledec's splitting

$T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

- 1 either *trivial* $T_\Lambda M = E^0$, i.e., all Lyapunov exponents at x are zero;
- 2 or *uniformly hyperbolic* with $E^u = E^+$, $E^s = E^-$;
- 3 or *partially hyperbolic* with $E^u = E^+$, $E^c = E^0$, $E^s = E^-$ (all $\neq \{0\}$).

Moreover, the 2nd alternative occurs for a positive μ -measure set of points $x \in M$ if and only if $f \in \mathcal{R}$ is Anosov.

Rem.: A weaker version (with no PH) was proved earlier in [B., Viana '05].

Discussion: $\dim M = 2$

If $\dim M = 2$ then the 3rd alternative in the theorem (partial hyperbolicity with 3 bundles) is impossible, so we get:

Corollary (B. '02)

C^1 -generic area-preserving diffeomorphisms are either Anosov or have zero Lyapunov exponents almost everywhere.

Rem.: The proof of this result appeared much before, and relied heavily on Mañé's ideas.

Discussion: $\dim M > 2$

For $\dim M > 2$, the picture is not necessarily so nice. . .

Discussion: $\dim M > 2$

For $\dim M > 2$, the picture is not necessarily so nice. . .

One can break M (minus a zero set) invariantly:

$$M = Z \sqcup \bigsqcup \Lambda_n \text{ mod } 0 \text{ where } \begin{cases} Z = \{\text{all } \lambda_i = 0\}, \\ \Lambda_n = \text{partially hyperbolic sets} \end{cases}$$

- Each Λ_n (or its closure $\overline{\Lambda}_n$) has of course its hyperbolicity constants c_n, τ_n .
- However, these constants become weaker and weaker as n grows.

Discussion: $\dim M > 2$

For $\dim M > 2$, the picture is not necessarily so nice. . .

One can break M (minus a zero set) invariantly:

$$M = Z \sqcup \bigsqcup \Lambda_n \text{ mod } 0 \text{ where } \begin{cases} Z = \{\text{all } \lambda_i = 0\}, \\ \Lambda_n = \text{partially hyperbolic sets} \end{cases}$$

- Each Λ_n (or its closure $\overline{\Lambda}_n$) has of course its hyperbolicity constants c_n, τ_n .
- However, these constants become weaker and weaker as n grows.

Please note that this is much stronger than what's is given by Oseledets theorem, which gives no uniformity along the orbits.

Also note that if $f \in \mathcal{R}$ is ergodic then the situation is much simpler. . .

The case of globally partially hyperbolic maps

Let $\text{PH}_\omega^1(M)$ be the subset of $\text{Diff}_\omega^1(M)$ formed by the diffeos that have a partially hyperbolic splitting over the whole tangent bundle.

The case of globally partially hyperbolic maps

Let $\text{PH}_\omega^1(M)$ be the subset of $\text{Diff}_\omega^1(M)$ formed by the diffeos that have a partially hyperbolic splitting over the whole tangent bundle.

Theorem (B. '10)

For the generic $f \in \text{PH}_\omega^1(M)$, all Lyapunov exponents in the center bundle are zero almost everywhere.

Rem: It may be necessary to pass to a different global partially hyperbolic splitting.

The case of globally partially hyperbolic maps

Let $\text{PH}_\omega^1(M)$ be the subset of $\text{Diff}_\omega^1(M)$ formed by the diffeos that have a partially hyperbolic splitting over the whole tangent bundle.

Theorem (B. '10)

For the generic $f \in \text{PH}_\omega^1(M)$, all Lyapunov exponents in the center bundle are zero almost everywhere.

Rem: It may be necessary to pass to a different global partially hyperbolic splitting.

Proof of the theorem: Just combine the previous theorem with this:

Theorem (Corollary of Dolgopyat–Wilkinson)

Generic $f \in \text{PH}_\omega^1(M)$ are (accessible and) weakly ergodic (i.e., almost every point has a dense orbit).

The case of globally partially hyperbolic maps

Let $\text{PH}_\omega^1(M)$ be the subset of $\text{Diff}_\omega^1(M)$ formed by the diffeos that have a partially hyperbolic splitting over the whole tangent bundle.

Theorem (B. '10)

For the generic $f \in \text{PH}_\omega^1(M)$, all Lyapunov exponents in the center bundle are zero almost everywhere.

Rem: It may be necessary to pass to a different global partially hyperbolic splitting.

Nice thing: The zero exponents in the center give a *nonuniform version of Burns–Wilkinson's center bunching*.

Ergodicity

Indeed many of Burns–Wilkinson's arguments work with *nonuniform center bunching*.

Putting these together with other [non obvious!] arguments, we get:

Theorem (Avila, B., Wilkinson '09)

The generic $f \in \text{PH}_\omega^1(M)$ is ergodic.

(**Curiosity:** This paper was published before its ancestors [BW'10] and [B'10].)

This gives a C^1 -generic, symplectic version of the Pugh–Shub ergodicity conjecture.

Proofs?

Now let's give an idea of the proof of the theorem stated by Mañé:

Theorem

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledecs splitting $T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

- 1 either *trivial* $T_\Lambda M = E^0$, i.e., all Lyapunov exponents at x are zero;
- 2 or *uniformly hyperbolic* with $E^u = E^+$, $E^s = E^-$;
- 3 or *partially hyperbolic* with $E^u = E^+$, $E^c = E^0$, $E^s = E^-$ (all $\neq \{0\}$).

Moreover, the 2nd alternative occurs for a positive μ -measure set of points $x \in M$ if and only if $f \in \mathcal{R}$ is Anosov.

Proofs?

Now let's give an idea of the proof of the theorem stated by Mañé:

Theorem

For every f in a residual (dense G_δ) subset \mathcal{R} of $\text{Diff}_\omega^1(M)$, the following properties hold: For μ -a.e. $x \in M$, the zipped Oseledets splitting $T_\Lambda M = E^+ \oplus E^0 \oplus E^-$ on $\Lambda = \text{orb}(x) = \{f^n(x); n \in \mathbb{Z}\}$ is:

- 1 either *trivial* $T_\Lambda M = E^0$, i.e., all Lyapunov exponents at x are zero;
- 2 or *uniformly hyperbolic* with $E^u = E^+$, $E^s = E^-$;
- 3 or *partially hyperbolic* with $E^u = E^+$, $E^c = E^0$, $E^s = E^-$ (all $\neq \{0\}$).

Moreover, the 2nd alternative occurs for a positive μ -measure set of points $x \in M$ if and only if $f \in \mathcal{R}$ is Anosov.

For the “moreover” part, we show that for C^2 (and hence C^1 -generic) diffeos, hyperbolic sets have either zero or full measure. [B., Viana '04].

Back to basics: domination, and the symplectic case

In fact, to prove the result above, one “only” needs to show the following:

Theorem (Main Theorem)

If f is generic in $\text{Diff}_\omega^1(M)$ then for almost every $x \in M$, the Oseledets splitting along the orbit of x is either trivial or dominated.

A Df -invariant splitting $T_\Lambda M = E^1 \oplus \dots \oplus E^k$ is called

- *trivial* if $k = 1$;
- *dominated* if each E_i dominates E_{i+1} .

Back to basics: domination, and the symplectic case

In fact, to prove the result above, one “only” needs to show the following:

Theorem (Main Theorem)

If f is generic in $\text{Diff}_\omega^1(M)$ then for almost every $x \in M$, the Oseledets splitting along the orbit of x is either trivial or dominated.

A Df -invariant splitting $T_\Lambda M = E^1 \oplus \dots \oplus E^k$ is called

- *trivial* if $k = 1$;
- *dominated* if each E_i dominates E_{i+1} .

Then, to conclude the proof of Mañé’s statement, one has to use that **for symplectic maps, “domination implies partially hyperbolicity”**. [B., Viana '04].

Rem.: Actually we obtain more information than in Mañé’s statement, since also get domination between different exponents of the same sign.

Comparison with the volume-preserving case

The “Main Theorem” just stated is also true replacing “symplectic” by “volume-preserving”:

Theorem (B. Viana '05)

If f is generic in $\text{Diff}_{\text{vol}}^1(M)$ then for almost every $x \in M$, the Oseledets splitting along the orbit of x is either trivial or dominated.

Comparison with the volume-preserving case

The “Main Theorem” just stated is also true replacing “symplectic” by “volume-preserving”:

Theorem (B. Viana '05)

If f is generic in $\text{Diff}_{\text{vol}}^1(M)$ then for almost every $x \in M$, the Oseledets splitting along the orbit of x is either trivial or dominated.

The proof of the two results have many things in common; however the symplectic result is more difficult, and we will see...

Comparison with the volume-preserving case

The “Main Theorem” just stated is also true replacing “symplectic” by “volume-preserving”:

Theorem (B. Viana '05)

If f is generic in $\text{Diff}_{\text{vol}}^1(M)$ then for almost every $x \in M$, the Oseledets splitting along the orbit of x is either trivial or dominated.

The proof of the two results have many things in common; however the symplectic result is more difficult, and we will see...

Rem.: There are some recent “global” improvements of this result: Avila–B. (ArXiv 2010), Jana Rodriguez–Hertz (in preparation), but that’s another story...

Another reduction

Integrated summed Lyapunov exponent:

$$\begin{aligned}L_p(f) &= \int_M (\lambda_1 + \cdots + \lambda_p) d\mu \\ &= \int \left(\lim_{n \rightarrow \infty} \log \| \wedge^p (Df^n) \| \right) d\mu.\end{aligned}$$

Easy fact: $L_p : \text{Diff}_\omega^1(M) \rightarrow \mathbb{R}$ is upper-semicontinuous, and thus continuous on a residual subset.

Another reduction

Integrated summed Lyapunov exponent:

$$\begin{aligned}L_p(f) &= \int_M (\lambda_1 + \cdots + \lambda_p) d\mu \\ &= \int \left(\lim_{n \rightarrow \infty} \log \| \wedge^p (Df^n) \| \right) d\mu.\end{aligned}$$

Easy fact: $L_p : \text{Diff}_\omega^1(M) \rightarrow \mathbb{R}$ is upper-semicontinuous, and thus continuous on a residual subset.

Let $N = d/2$.

Theorem

If f is a point of continuity of L_1, \dots and L_N then the Oseledets splitting is trivial or dominated along almost every orbit.

Another reduction

Integrated summed Lyapunov exponent:

$$\begin{aligned}L_p(f) &= \int_M (\lambda_1 + \cdots + \lambda_p) d\mu \\ &= \int \left(\lim_{n \rightarrow \infty} \log \| \wedge^p (Df^n) \| \right) d\mu.\end{aligned}$$

Easy fact: $L_p : \text{Diff}_\omega^1(M) \rightarrow \mathbb{R}$ is upper-semicontinuous, and thus continuous on a residual subset.

Let $N = d/2$.

Theorem

If f is a point of continuity of L_1, \dots and L_N then the Oseledets splitting is trivial or dominated along almost every orbit.

Strategy of the proof: Suppose that one can detect non-domination of the Oseledets splitting on a positive measure set. Then produce a perturbation of f for which some L_p drops.

Setup for the proof

Fix f , $p \in \{1, \dots, N = d/2\}$. Assume that the following set has positive measure:

$$\Sigma_p = \{x \in M; x \text{ regular non periodic with } \lambda_p(x) > \lambda_{p+1}(x)\}$$


Setup for the proof

Fix f , $p \in \{1, \dots, N = d/2\}$. Assume that the following set has positive measure:

$$\Sigma_p = \{x \in M; x \text{ regular non periodic with } \lambda_p(x) > \lambda_{p+1}(x)\}$$

Another “zipped” Oseledets splitting:

$$T_{\Sigma_p} M = E^u \oplus E^c \oplus E^s \quad \text{where} \begin{cases} \dim E^u = \dim E^s = p \\ \dim E^c = 2(N - p) \end{cases}$$

 Misleading notation: the splitting is not PH!


Setup for the proof

Fix f , $p \in \{1, \dots, N = d/2\}$. Assume that the following set has positive measure:

$$\Sigma_p = \{x \in M; x \text{ regular non periodic with } \lambda_p(x) > \lambda_{p+1}(x)\}$$

Another “zipped” Oseledets splitting:

$$T_{\Sigma_p} M = E^u \oplus E^c \oplus E^s \quad \text{where} \begin{cases} \dim E^u = \dim E^s = p \\ \dim E^c = 2(N - p) \end{cases}$$

 Misleading notation: the splitting is not PH!

Denote $E^{uc} = E^u \oplus E^c$ etc. Then

$$\omega(E^u, E^{uc}) \equiv \omega(E^c, E^{us}) \equiv \omega(E^s, E^{cs}) \equiv 0.$$


Setup for the proof

Fix f , $p \in \{1, \dots, N = d/2\}$. Assume that the following set has positive measure:

$$\Sigma_p = \{x \in M; x \text{ regular non periodic with } \lambda_p(x) > \lambda_{p+1}(x)\}$$

Another “zipped” Oseledets splitting:

$$T_{\Sigma_p} M = E^u \oplus E^c \oplus E^s \quad \text{where} \begin{cases} \dim E^u = \dim E^s = p \\ \dim E^c = 2(N - p) \end{cases}$$

 Misleading notation: the splitting is not PH!

Denote $E^{uc} = E^u \oplus E^c$ etc. Then

$$\omega(E^u, E^{uc}) \equiv \omega(E^c, E^{us}) \equiv \omega(E^s, E^{cs}) \equiv 0.$$

Strategy: Assume $E^u \not\perp E^{cs}$ and perturb f so that $L_p = \int(\lambda_1 + \dots + \lambda_p)$ drops.

Main steps (sketchy)

- If a point in Σ_p “sees” non-domination $E^u \not\prec E^{cs}$ (for example, if $\angle(E^u, E^{cs})$ is small) then we can find a perturbation g of f that sends a vector from E_f^u to E_f^{cs} .

Main steps (sketchy)

- If a point in Σ_p “sees” non-domination $E^u \not\asymp E^{cs}$ (for example, if $\angle(E^u, E^{cs})$ is small) then we can find a perturbation g of f that sends a vector from E_f^u to E_f^{cs} .
- Do that in the *middle* of a long segment of orbit $\{x, \dots, f^n x\}$. Then one gets

$$\frac{1}{n} \log \|\wedge^p Dg^n(x)\| < \lambda_1(f, x) + \dots + \lambda_p(f, x) - \text{Something.}$$

Main steps (sketchy)

- If a point in Σ_p “sees” non-domination $E^u \not\prec E^{cs}$ (for example, if $\angle(E^u, E^{cs})$ is small) then we can find a perturbation g of f that sends a vector from E_f^u to E_f^{cs} .
- Do that in the *middle* of a long segment of orbit $\{x, \dots, f^n x\}$. Then one gets

$$\frac{1}{n} \log \|\wedge^p Dg^n(x)\| < \lambda_1(f, x) + \dots + \lambda_p(f, x) - \text{Something.}$$

- ▶ Something = $\frac{1}{2}(\lambda_p(f, x) + \lambda_{p+1}(f, x))$ (which is POSITIVE).
- ▶ In fact we should obtain the inequality not only for x [a zero measure set is useless], but for most z around x in the support of the perturbation.

Main steps (continued)

Example: $\dim M = 2$

$$Df^n(x) = \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix}^k \text{Id}^m \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix}^k, \quad k \simeq n/2 \gg m \gg 1.$$

[draw a figure with the solution]

Main steps (continued)

Example: $\dim M = 2$

$$Df^n(x) = \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix}^k \text{Id}^m \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix}^k, \quad k \simeq n/2 \gg m \gg 1.$$

[draw a figure with the solution]

Back to the steps of the general proof:

- Around a segment of orbit $\{x, \dots, f^n x\}$ that sees nondomination we find a thin and long tower $U \sqcup f(U) \sqcup \dots \sqcup f^n(U)$, and find a perturbation supported in the tower so that the “finite-time” summed exponent drops (as explained above).
- Cover the a (“large”) positive measure set of the manifold with these towers.
- This causes a significant drop of $L_p = \int (\lambda_1 + \dots + \lambda_p)$, as we wanted.

The 4 types of non-dominance

Assume $x \in \Sigma_p$ and the segment of orbit $\{x, \dots, f^m x\}$ is very long ($m \gg 1$) and “sees” non-domination $E^u \not\asymp E^{cs}$; more precisely:

$$\frac{\|Df^m(x)|E^{cs}(x)\|}{m(Df^m(x)|E^u(x))} \geq \frac{1}{2}. \quad (m(L) = \|L^{-1}\|^{-1}).$$

Lemma (of Symplectic Linear Algebra): one of the following 4 cases occurs:

The 4 types of non-dominance

Assume $x \in \Sigma_p$ and the segment of orbit $\{x, \dots, f^m x\}$ is very long ($m \gg 1$) and “sees” non-domination $E^u \not\asymp E^{cs}$; more precisely:

$$\frac{\|Df^m(x)|E^{cs}(x)\|}{m(Df^m(x)|E^u(x))} \geq \frac{1}{2}. \quad (m(L) = \|L^{-1}\|^{-1}).$$

Lemma (of Symplectic Linear Algebra): one of the following 4 cases occurs:

Case I: small angle. There is a point $y \in \{x, \dots, f^m x\}$ such that $\angle(E^{cs}(y), E^u(y)) \ll 1$.

How to perturb it: Compose with a single small rotation.

The 4 types of non-dominance

Assume $x \in \Sigma_p$ and the segment of orbit $\{x, \dots, f^m x\}$ is very long ($m \gg 1$) and “sees” non-domination $E^u \not\asymp E^{cs}$; more precisely:

$$\frac{\|Df^m(x)|E^{cs}(x)\|}{m(Df^m(x)|E^u(x))} \geq \frac{1}{2}. \quad (m(L) = \|L^{-1}\|^{-1}).$$

Lemma (of Symplectic Linear Algebra): one of the following 4 cases occurs:

Case I: small angle. There is a point $y \in \{x, \dots, f^m x\}$ such that $\angle(E^{cs}(y), E^u(y)) \ll 1$.

How to perturb it: Compose with a single small rotation.

Case II: inverted behaviors. There are unit vectors $v^{cs} \in E^{cs}(x)$, $v^u \in E^u(x)$ such that $\|Df^m(x) \cdot v^{cs}\| \gg \|Df^m(x) \cdot v^u\|$. (Replace if necessary the whole segment of orbit by a subsegment.)

How to perturb it: Use two small rotations, one at the beginning and the other at the end.

The 4 types of non-dominance (continued)

Case III: Identity on a symplectic ($\omega \neq 0$) plane.

There is a long subsegment of orbit such the following holds: There is a (2-dim) plane spanned by a vector in E^u and a vector in E^s such that the restriction of Df to this plane “looks like the identity” (or more precisely, becomes an isometry after a bounded change of coordinates).

Notice that $\omega \neq 0$ in P .

The 4 types of non-dominance (continued)

Case III: Identity on a symplectic ($\omega \neq 0$) plane.

There is a long subsegment of orbit such the following holds: There is a (2-dim) plane spanned by a vector in E^u and a vector in E^s such that the restriction of Df to this plane “looks like the identity” (or more precisely, becomes an isometry after a bounded change of coordinates).

Notice that $\omega \neq 0$ in P .

How to perturb it: This case is essentially 2-dimensional. Use several “nested” rotations, as explained in a previous example. . .

The 4 types of non-dominance (continued)

Case IV: Expansion on a null ($\omega \equiv 0$) plane.

There is a (2-dim) plane P spanned by a vector in E^u and a vector in E^c which is **uniformly expanded** and **conformal** (along the segment of orbit). That is, after a bounded change of coordinates we have

$$Df(f^i x)|_P = \begin{pmatrix} \tau_i & 0 \\ 0 & \tau_i \end{pmatrix}, \quad \tau_i > c > 1.$$

The plane is necessarily null ($\omega \equiv 0$).

How to perturb it?

The 4 types of non-dominance (continued)

Case IV: Expansion on a null ($\omega \equiv 0$) plane.

There is a (2-dim) plane P spanned by a vector in E^u and a vector in E^c which is **uniformly expanded** and **conformal** (along the segment of orbit). That is, after a bounded change of coordinates we have

$$Df(f^i x)|_P = \begin{pmatrix} \tau_i & 0 \\ 0 & \tau_i \end{pmatrix}, \quad \tau_i > c > 1.$$

The plane is necessarily null ($\omega \equiv 0$).

How to perturb it?

The 1st idea would be to imitate the previous case: rotate the plane P around a complementary codimension-2 axis (where the rotation is the identity).

The 4 types of non-dominance (continued)

Case IV: Expansion on a null ($\omega \equiv 0$) plane.

There is a (2-dim) plane P spanned by a vector in E^u and a vector in E^c which is **uniformly expanded** and **conformal** (along the segment of orbit). That is, after a bounded change of coordinates we have

$$Df(f^i x)|_P = \begin{pmatrix} \tau_i & 0 \\ 0 & \tau_i \end{pmatrix}, \quad \tau_i > c > 1.$$

The plane is necessarily null ($\omega \equiv 0$).

How to perturb it?

The 1st idea would be to imitate the previous case: rotate the plane P around a complementary codimension-2 axis (where the rotation is the identity).

However, this perturbation is not symplectic!

(In the **volume-preserving** case this idea would work; there are only 3 cases to be considered there.)

4-dimensional problem

There are standard symplectic coordinates $p_1, \dots, p_N, q_1, \dots, q_N$ (so $\omega = \sum_i dp_i \wedge dq_i$) such that

$$P = \left\langle \underbrace{\frac{\partial}{\partial p_1}}_{\in E^u}, \underbrace{\frac{\partial}{\partial p_2}}_{\in E^c} \right\rangle,$$

$$Q = \left\langle \underbrace{\frac{\partial}{\partial q_2}}_{\in E^c}, \underbrace{\frac{\partial}{\partial p_1}}_{\in E^s} \right\rangle.$$

$$Df|_{P \oplus Q} = \begin{pmatrix} \tau_i & & & \\ & \tau_i & & \\ & & \tau_i^{-1} & \\ & & & \tau_i^{-1} \end{pmatrix} \quad (\text{order: } p_1, p_2, q_2, q_1.)$$

If we rotate P we also need to rotate Q .

4-dimensional problem

There are standard symplectic coordinates $p_1, \dots, p_N, q_1, \dots, q_N$ (so $\omega = \sum_i dp_i \wedge dq_i$) such that

$$P = \left\langle \underbrace{\frac{\partial}{\partial p_1}}_{\in E^u}, \underbrace{\frac{\partial}{\partial p_2}}_{\in E^c} \right\rangle,$$
$$Q = \left\langle \underbrace{\frac{\partial}{\partial q_2}}_{\in E^c}, \underbrace{\frac{\partial}{\partial p_1}}_{\in E^s} \right\rangle.$$

$$Df|_{P \oplus Q} = \begin{pmatrix} \tau_i & & & \\ & \tau_i & & \\ & & \tau_i^{-1} & \\ & & & \tau_i^{-1} \end{pmatrix} \quad (\text{order: } p_1, p_2, q_2, q_1.)$$

If we rotate P we also need to rotate Q .

Nested rotations don't work! The problem is that hyperbolicity of Df quickly distorts the domain where the perturbation should be supported.

Solution of Case IV

- Start with a box D as a perturbation domain. (We can pretend $M = \mathbb{R}^4$.)

Solution of Case IV


- Start with a box D as a perturbation domain. (We can pretend $M = \mathbb{R}^4$.)
- Choose **any** symplectic perturbation of the identity $h: D \rightarrow D$ that *doesn't* leave the field of directions $\frac{\partial}{\partial \rho_1}$ invariant. Let $g = f \circ h$ be the perturbation of f .

Solution of Case IV

- Start with a box D as a perturbation domain. (We can pretend $M = \mathbb{R}^4$.)
- Choose **any** symplectic perturbation of the identity $h: D \rightarrow D$ that *doesn't* leave the field of directions $\frac{\partial}{\partial p_1}$ invariant. Let $g = f \circ h$ be the perturbation of f .
- Let Θ_0 be the angle (in the $p_1 p_2$ projection) that the field of directions $v_0 = \frac{\partial}{\partial p_1}$ is rotated as we apply h . View Θ_0 as a **random variable**. (Normalize measure $\mu(D) = 1$).

Solution of Case IV

- Start with a box D as a perturbation domain. (We can pretend $M = \mathbb{R}^4$.)
- Choose **any** symplectic perturbation of the identity $h: D \rightarrow D$ that *doesn't* leave the field of directions $\frac{\partial}{\partial p_1}$ invariant. Let $g = f \circ h$ be the perturbation of f .
- Let Θ_0 be the angle (in the $p_1 p_2$ projection) that the field of directions $v_0 = \frac{\partial}{\partial p_1}$ is rotated as we apply h . View Θ_0 as a **random variable**. (Normalize measure $\mu(D) = 1$).
- Look the image $g(D)$ and the image v_1 of the field $v_0 = \frac{\partial}{\partial p_1}$ by Dg . Using Vitali Lemma, cover most of $g(D)$ by many disjoint tiny boxes D_i that look like (basically after a change of scale) the original box:



Solution of Case IV (continued)

- Inside each small box D_i , we make a perturbation which is just a rescaling of h .

Solution of Case IV (continued)

- Inside each small box D_i , we make a perturbation which is just a rescaling of h .
- Let Θ_1 be the angle (in the $p_1 p_2$ projection) that v_1 is rotated at this step. We view Θ_1 as a random variable ($\mu(g(D)) = 1$).

Solution of Case IV (continued)

- Inside each small box D_i , we make a perturbation which is just a rescaling of h .
- Let Θ_1 be the angle (in the p_1p_2 projection) that v_1 is rotated at this step. We view Θ_1 as a random variable ($\mu(g(D)) = 1$).
- Then **THE RANDOM VARIABLES Θ_0 AND Θ_1 ARE (approximately) INDEPENDENT AND IDENTICALLY DISTRIBUTED!**

Solution of Case IV (continued)

- Inside each small box D_i , we make a perturbation which is just a rescaling of h .
- Let Θ_1 be the angle (in the $p_1 p_2$ projection) that v_1 is rotated at this step. We view Θ_1 as a random variable ($\mu(g(D)) = 1$).
- Then **THE RANDOM VARIABLES Θ_0 AND Θ_1 ARE (approximately) INDEPENDENT AND IDENTICALLY DISTRIBUTED!**
- Continuing this process we obtain a **RANDOM WALK**
 $S_n = \Theta_0 + \Theta_1 + \cdots + \Theta_n$.
- Since every (non-stopped) random walk is transient, most orbits eventually reach $\pm\pi/2$.
- Thus we succeeded in sending the direction $\frac{\partial}{\partial p_1} \in E^u$ to the direction $\frac{\partial}{\partial q_1} \in E^c$ (for most points).

The end

THANK YOU!